

**References for Joslin's Clinical Nutrition Guideline for Overweight and
Obese Adults with Type 2 Diabetes, Prediabetes or Those at High Risk for
Developing Type 2 Diabetes**
3/29/07

Target population

1. Deurenberg P, Yap M, Staveren WA. Body mass index and percent body fat: a meta-analysis among different ethnic groups. *Int J Obes* 22:1164-1171, 1998.
2. Eriksson KF, Lindgarde F. Prevention of type 2 (non-insulin dependent) diabetes mellitus by diet and physical exercise. The 6-year Malmo feasibility study. *Diabetologia* 34:891-8, 1991.
3. Grundy SM, Cleeman JI, Daniels SR et al. Diagnosis and management of the metabolic syndrome. An American Heart Association/National Heart, Lung, and Blood Institute Scientific Statement. *Circulation* 112:2735-52, 2005.
4. Hunking P. Lifestyle advice in preventing and treating type 2 diabetes. *J Fam Health Care* 16:109-12, 2006.
5. Misra A, Wasir JS, Pandey RM. An evaluation of candidate definitions of the metabolic syndrome in adult Asian Indians. *Diabetes Care* 28:398-403, 2005.

General Recommendations

1. Cernea S, Hancu N, Raz I. Diet and coronary heart disease in diabetes. *Acta Diabetol* 40:S389-400, 2003.
2. Gaetke LM, Stuart MA, Trusczyńska H. A single nutrition counseling session with a registered dietitian improves short-term clinical outcomes for rural Kentucky patients with chronic diseases. *J Am Diet Assoc* 106:109-12, 2006.
3. Parillo M, Riccardi G. Diet composition and the risk of type 2 diabetes: epidemiological and clinical evidence. *Br J Nutr* 92:7-19, 2004.

Weight Reduction

1. Beach J. Winning at weight loss. The drug option. Weight-loss medications can help, but they are not side effect free. *Diabetes Forecast* 59:55-7, April 2006.
2. Brown S., Upchurch S., Anding R., Winter M., Ramirez G. Promoting Weight Loss in Type II Diabetes. *Diabetes Care* 19:613-624, 1996.

3. Dansinger ML, Gleason JA, Griffith JL et al. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA* 293:43-53, 2005.
4. Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Eng J Med* 346:393-403, 2002.
5. Dixon JB, O'Brien PE. Health outcomes of severely obese type 2 diabetic subjects 1 year after laparoscopic adjustable gastric banding. *Diabetes Care* 25:358-63, 2002.
6. Dixon JB. Weight loss medications: where do they fit in? *Aus Fam Physician* 35:576-9, 2006.
7. Ferchak CV, Meneghini LF. Obesity, bariatric surgery and type 2 diabetes--a systematic review. *Diabetes Metab Res Rev* 20:438-45, 2004.
8. Hamdy O, Ledbury S, Mullooly C et al. Lifestyle modification improves endothelial function in obese subjects with the insulin resistance syndrome. *Diabetes Care* 26:2119-25, 2003.
9. Hamman RF et al. Effect of weight loss with lifestyle intervention on risk of diabetes. *Diabetes Care* 29:2102-7, 2006.
10. Kosaka K, Noda M, Kuzuya T. Prevention of type 2 diabetes by lifestyle intervention: a Japanese trial in IGT males. *Diabetes Res Clin Pract* 67:152-62, 2005.
11. Lara-Castro C, Garvey T. Diet, insulin resistance, and obesity: zoning in on data for Atkins dieters living in South Beach. *J Clin Endocrinol Metab* 89:4197-4205, 2004.
12. Liz Z, Hong K, Saltsman P et al. Long-term efficacy of soy-based meal replacements vs an individualized diet plan in obese type II DM patients: relative effects on weight loss, metabolic parameters and C-reactive protein. *Eur J Clin Nutr* 59:411-8, 2005.
13. Li Z, Maglione M, Tu W et al. Meta-analysis: pharmacologic treatment of obesity. *Ann Intern Med* 5; 142:532-46, 2005.
14. Mann JJ. Diet and risk of coronary heart disease and type 2 diabetes. *Lancet* 360:783-9, 2002.
15. Moore H, Summerbell C, Hooper L et al. Dietary advice for treatment of type 2 diabetes mellitus in adults. *Cochrane Database Syst Rev* (3):CD004097, 2004.

16. Noakes M, Foster PR, Keogh JB, Clifton PM. Are meal replacements an effective strategy for treating obesity in adults with features of metabolic syndrome? *Asia Pac J Clin Nutr* 13:S63, 2004.
17. Noakes M, Foster PR, Keogh JB, Clifton PM. Meal replacements are as effective as structured weight-loss diets for treating obesity in adults with features of metabolic syndrome. *J Nutr* 134:1894-9, 2004.
18. Norris SL, Xuanping Z, Avenell A et al. Long-term effectiveness of lifestyle and behavioral weight loss interventions in adults with type 2 diabetes: a meta-analysis. *Am J Med* 117:762-774, 2004.
19. Steyn NP, Mann J, Bennett PH et al. Diet, nutrition and the prevention of type 2 diabetes. *Public Health Nutr* 7:147-65, 2004.
20. Torgerson JS, Hauptman J, Boldrin MN, Sjostrom L. XENical in the prevention of diabetes in obese subjects (XENDOS) study: a randomized study of orlistat as an adjunct to lifestyle changes for the prevention of type 2 diabetes in obese patients. *Diabetes Care* 27:155-61, 2004.
21. Uusitupa M, Lindi V, Louheranta A et al. Finnish Diabetes Prevention Study Group. Long-term improvement in insulin sensitivity by changing lifestyles of people with impaired glucose tolerance: 4-year results from the Finnish Diabetes Prevention Study. *Diabetes* 52:2532-8, 2003.
22. Wadden TA, Berkowitz RI, Womble LG et al. Randomized trial of lifestyle modification and pharmacotherapy for obesity. *N Engl J Med* 353:2111-20, 2005.
23. Yamaoka K, Tango T. Efficacy of lifestyle education to prevent type 2 diabetes. *Diabetes Care* 28:2780-2786, 2005.
24. Yip I, Go VL, DeShields S et al. Liquid meal replacements and glycemic control in obese type 2 diabetes patients. *Obes Res* 9:341S-347S, 2001.

Carbohydrate

1. Anderson JW, Allgood LD, Turner J et al. Effects of psyllium on glucose and serum lipid responses in men with type 2 diabetes and hypercholesterolemia. *Am J Clin Nutr* 70:466-73, 1999.
2. Anderson JW, Gustafson NJ, Bryant CA, Tietyan-Clark J. Dietary fiber and diabetes: A comprehensive review and practical application. *J Am Diet Assoc* 87:1189-97, 1987.

3. Baba NH, Sawaya S, Torbay N et al. High protein vs high carbohydrate hypoenergetic diet for the treatment of obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord* 23:1202-1206, 1999.
4. Biesalski HK. Diabetes preventive components in the Mediterranean diet. *Eur J Nutr* 43:I/26-30, 2004.
5. Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA. A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Clin Endocrinol Metab* 88:1617-23, 2003.
6. Brinkworth GD, Noakes M, Keogh JB et al. Long-term effects of a high-protein, low-carbohydrate diet on weight control and cardiovascular risk markers in obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord* 28:661-70, 2004.
7. Chandalia M, Garg A, Lutjohann D et al. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *N Engl J Med* 342:1392-1398, 2000.
8. Dumesnil JG, Turgeon J, Tremblay A et al. Effect of a low-glycaemic index--low-fat--high protein diet on the atherogenic metabolic risk profile of abdominally obese men. *Br J Nutr* 86:557-68, 2001.
9. Foster GD, Wyatt HR, Hill JO et al. A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med* 348:2082-90, 2003.
10. Fukagawa NK, Anderson JW, Hageman G et al. High-carbohydrate, high-fiber diets increase peripheral insulin sensitivity in healthy young and old adults. *Am J Clin Nutr* 52:524-8, 1990.
11. Gannon MC, Nuttall FQ, Neil BJ, Westphal SA. The insulin and glucose responses to meals of glucose plus various proteins in type II diabetic subjects. *Metabolism* 37:1081-8, 1988.
12. Heilbronn LK, Noakes M, Clifton PM. The effect of high- and low-glycemic index energy restricted diets on plasma lipid and glucose profiles in type 2 diabetic subjects with varying glycemic control. *J Am Coll Nutr* 21:120-7, 2002.
13. Hu FB. Plant-based foods and prevention of cardiovascular disease: an overview. *Am J Clin Nutr* 78:544S-551S, 2003.
14. Keno Y, Matsuzawa Y, Tokunaga K et al. High sucrose diet increases visceral fat accumulation in VMH-lesioned obese rats. *Int J Obes* 15:205-11, 1991.

15. Liu S, Willett WC, Manson JE et al. Relation between changes in intakes of dietary fiber and grain products and changes in weight and development of obesity among middle-aged women. *Am J Clin Nutr* 78:920-7, 2003.
16. Ludwig DS, Pereira MA, Kroenke CH et al. Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA* 282:1539-46, 1999.
17. Ludwig DS. Dietary glycemic index and obesity. *J Nutr* 130:280S-3S, 2000.
18. Miyashita Y, Koide N, Ohtsuka M et al. Beneficial effect of low carbohydrate in low calorie diets on visceral fat reduction in type 2 diabetic patients with obesity. *Diabetes Res Clin Pract* 65:235-41, 2004.
19. Pereira MA, Swain J, Goldfine AB et al. Effects of a low glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA* 292:2482-90, 2004.
20. Stern L, Iqbal N, Seshadri P et al. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med* 140:778-85, 2004.
21. Story L, Anderson JW, Chen WJ et al. Adherence to high-carbohydrate, high-fiber diets: Long-term studies of non-obese diabetic men. *J Am Diet Assoc* 85:1105-10, 1985.
22. Westman EC, Yancy WS, Edman JS et al. Effect of 6-month adherence to a very low carbohydrate diet program. *Am J Med* 113:30-6, 2002.
23. Ziai SA. Psyllium decreased serum glucose and glycosylated hemoglobin significantly in diabetic outpatients. *J Ethnopharmacol* 102:202-7, 2005.

Fat

1. Barnard ND, Cohen J, Jenkins DJ et al. A low-fat vegan diet improves glycemic control and cardiovascular risk factors in a randomized clinical trial in individuals with type 2 diabetes. *Diabetes Care* 29:1777-1783, 2006.
2. Fleming RM. The effect of high-, moderate-, and low-fat diets on weight loss and cardiovascular disease risk factors. *Prev Cardiol* 5:110-8, 2002.

3. Hodson L, Skeaff CM, Chisholm WA. The effect of replacing dietary saturated fat with polyunsaturated or monounsaturated fat on plasma lipids in free-living young adults. *Eur J Clin Nutr* 55:908-15, 2001.
4. Hu FB, Stampfer MJ, Manson JE et al. Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med* 337:1491-9, 1997.
5. Julius U. Fat modification in the diabetes diet. *Exp Clin Endocrinol Diabetes* 111:60-5, 2003.
6. Kim JY, Nolte LA, Hansen PA et al. High-fat diet-induced muscle insulin resistance: relationship to visceral fat mass. *Am J Physiol Regul Integr Comp Physiol* 279:R2057-65, 2000.
7. McAuley KA, Hopkins CM, Smith KJ et al. Comparison of high-fat and high-protein diets with a high-carbohydrate diet in insulin-resistant obese women. *Diabetologia* 48:8-16, 2005.
8. Meckling KA, O'Sullivan C, Saari D. Comparison of a low-fat diet to a low-carbohydrate diet on weight loss, body composition, and risk factors for diabetes and cardiovascular disease in free-living, overweight men and women. *J Clin Endocrinol Metab* 89:2717-23, 2004.
9. Nappo F, Esposito K, Cioffi M et al. Postprandial endothelial activation in healthy subjects and in type 2 diabetic patients: role of fat and carbohydrate meals. *J Am Coll Cardiol* 39:1145-50, 2002.
10. Odegaard A, Pereira MA. Trans fatty acids, insulin resistance and type 2 diabetes. *Nutr Rev* 64:364-372, 2006.
11. Poppitt SD, Keogh GF, Mulvey TB et al. Effect of moderate changes in dietary fatty acid profile on postprandial lipaemia, haemostatic and related CVD risk factors in healthy men. *Eur J Clin Nutr* 58:819-27, 2004.
12. Rosenfalck AM, Almdal T, Viggers L et al. A low fat diet improves peripheral insulin sensitivity in patients with type 1 diabetes. *Diabetic Medicine* 23:384-86, 2006.
13. Salmeron J, Hu FB, Manson JE et al. Dietary fat intake and risk of type 2 diabetes in women. *Am J Clin Nutr* 73:1019-26, 2001.
14. Simopoulos AP. The importance of the ratio of omega-6/omega-3 essential fatty acids. *Biomed Pharmacother* 56:365-79, 2002.

15. Suzukawa M, Abbey M, Howe PR, Nestel PJ. Effects of fish oil fatty acids on low density lipoprotein size, oxidizability, and uptake by macrophages. *J Lipid Res* 36:473-84, 1995.
16. Mozaffarian D, Katan MB, Ascherio A, et al. Trans fatty acids and cardiovascular disease. *N Eng J Med* 354:1601-13, 2006.
17. Williams MJ, Sutherland WH, McCormick MP et al. Impaired endothelial function following a meal rich in used cooking fat. *J Am Coll Cardiol* 33:1050-5, 1999.

Protein

1. Brinkworth GD, Noakes M, Parker B et al. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia* 47:1677-86, 2004.
2. Gannon MC, Nuttall FQ, Saeed A et al. An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *Am J Clin Nutr* 78:734-41, 2003.
3. Gannon MC, Nuttall FQ. Effect of a high-protein, low-carbohydrate diet on blood glucose control in people with type 2 diabetes. *Diabetes* 53:2375-82, 2004.
4. Johnston CS, Tjonn SL, Swan PD. High-protein, low-fat diets are effective for weight loss and favorably alter biomarkers in healthy adults. *J Nutr* 134:586-91, 2004.
5. Lungershausen YK, Abbey M, Nestel PJ, Howe PR. Reduction of blood pressure and plasma triglycerides by omega-3 fatty acids in treated hypertensives. *J Hypertens* 12:1041-5, 1994.
6. Luscombe ND, Clifton PM, Noakes M et al. Effects of energy-restricted diets containing increased protein on weight loss, resting energy expenditure, and the thermic effect of feeding in type 2 diabetes. *Diabetes Care* 25:652-7, 2002.
7. Nuttall FQ, Gannon MC. Metabolic response of people with type 2 diabetes to a high protein diet. *Nutr Metab* 1:6, 2004.
8. Nuttall FQ, Gannon MC, Saeed A et al. The metabolic response of subjects with type 2 diabetes to a high-protein, weight-maintenance diet. *J Clin Endocrinol Metab* 88:3577-83, 2003.

9. Parker B, Noakes M, Luscombe N, Clifton P. Effect of a high-protein, high-monounsaturated fat weight loss diet on glycemic control and lipid levels in type 2 diabetes. *Diabetes Care* 25:425-30, 2002.
10. Sargrad KR, Homko C, Mozzoli M, Boden G. Effect of high protein vs high carbohydrate intake on insulin sensitivity, body weight, hemoglobin A1c, and blood pressure in patients with type 2 diabetes mellitus. *J Am Diet Assoc* 105:573-80, 2005.
11. Skov AR, Toubro S, Ronn B et al. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord* 23:528-36, 1999.
12. Wolfe BM, Piche LA. Replacement of carbohydrate by protein in a conventional-fat diet reduces cholesterol and triglyceride concentrations in healthy normolipidemic subjects. *Clin Invest Med* 22:140-8, 1999.

Physical Activity and Behavioral Modification

1. Normand GB, Haddad E, Kenny GP et al. Effects of exercise on glycemic control and body mass in type 2 diabetes mellitus. *JAMA* 286:1218-1227, 2001.
2. Pavlou KN, Krey S, Steffee WP. Exercise as an adjunct to weight loss and maintenance in moderately obese subjects. *Am J Clin Nutr* 49:1115-23, 1989.
3. Sigal RJ, Kenny GP, Wasserman DH et al. Physical activity/exercise and type 2 Diabetes: a consensus statement from the American Diabetes Association. *Diabetes Care* 29:1433-8, 2006.